

CMB CONSTRAINTS ON A BARYONIC DARK MATTER-DOMINATED UNIVERSE

LOUISE M GRIFFITHS, ALESSANDRO MELCHIORRI, JOSEPH SILK

Nuclear and Astrophysics Laboratory, University of Oxford,
 Keble Road, Oxford, OX 3RH, UK

Draft version February 1, 2008

ABSTRACT

In recent years, upper limits on the cosmic microwave background (CMB) anisotropies combined with predictions made by theories of galaxy formation, have been extremely powerful in ruling out purely baryonic dark matter (BDM) universes. However, it has recently been argued that the absence of a prominent second peak in the anisotropy spectrum measured by the BOOMERanG-98 and MAXIMA-1 experiments seems to favour a Λ BDM model when combined with standard Big Bang Nucleosynthesis (BBN) constraints. In this *Letter*, we further investigate this result showing that, using the CMB data *alone*, a purely baryonic adiabatic model of structure formation seems unlikely if the universe is flat ($\Omega = 1$). Combining the CMB data with supernova type Ia (SNIa) data renders purely baryonic models inconsistent with flatness at high significance and more than 3σ away from both the BBN constraints and the HST key project result $h = 0.72 \pm 0.08$. These results indicate that only a radical revision of cosmology with *ad hoc* properties could bring baryonic models such as those advocated by MOND enthusiasts back into agreement with current observations.

Subject headings: cosmology: cosmic microwave background — cosmology: dark matter

1. INTRODUCTION

Since the pioneering work of Zwicky (1933), astronomers have found compelling evidence to suggest that the major contribution to the overall mass density of the universe is in the form of *dark* matter. Even if its precise nature remains unknown (see e.g. Bergstrom 2000 for a recent review), the currently favoured hypothesis of *cold* dark matter (CDM) is in good agreement with an enormous body of data: anisotropies in the cosmic microwave background (CMB), large-scale structure surveys, cluster abundances, structure of the Lyman- α forest and a host of other measurements (see e.g. Bahcall et al. 1999). Furthermore, in the past 30 years, upper limits on the CMB anisotropies and predictions made by theories of galaxy formation have been extremely effective at ruling out models based on purely *baryonic* or *hot* dark matter (see e.g. Doroshkevich, Zel'dovich & Sunyaev 1978; Wilson & Silk 1981; Kaiser 1983; White, Frenk & Davis 1983).

The quality of astrophysical data has improved and it currently seems likely that the present fractional overall energy density Ω includes contributions from a cosmological constant Λ (Ω_Λ) and from cold (Ω_{cdm}), hot (Ω_{hdm}) and baryonic (Ω_b) dark matter. Also, it appears that the densities of these components are all within one or two orders of magnitude of each other. While the similar densities of dark ‘energy’ and ‘matter’ can possibly be explained by advocating a quintessence scalar field which ‘tracks’ the matter density (Steinhardt, Wang & Zlatev 1999), the reason why the densities of the 3 matter components are so similar is not so apparent.

Predictions made by structure evolution theories based on the Λ CDM model disagree with some galaxy observations, which suggests that, for the model to hold, assumptions made about the properties of CDM particles must be modified (see e.g. Spergel & Steinhardt 1999; Burkert 2000; Madsen 2000; Moore et al. 2000). If this

reasoning is adopted, currently discussed versions of the present cosmological scenario, incorporating a fine-tuned cosmological constant and dark matter particles with *ad hoc* properties, lose the compelling aesthetic simplicity of the original CDM model (see Sellwood & Kosowsky 2000 for a recent review). Of course, other, more astrophysical, resolutions may be possible Binney, Gerhard and Silk 2001.

The recent CMB anisotropy measurements from the BOOMERanG-98 (de Bernardis et al. 2000) and MAXIMA-1 (Hanany et al. 2000) experiments have provided new insights into the cosmological parameters. For the first time, the anisotropy angular power spectrum has been measured over a wide range of angular scales, from multipole $l \sim 50$ up to $l \sim 800$ with errors of the order of 10%. The data sets confirm that there is a peak in the angular power spectrum at $l \sim 200$ with a steep decline in power from $l \sim 200$ to $l \sim 300$. While the presence of such a peak is consistent with the predicted acoustic oscillations in the adiabatic inflationary scenario, the absence of prominent secondary peaks after $l \sim 300$ is an unexpected result, suggesting that the value for the physical baryonic density is $\sim 50\%$ higher than that predicted by Big Bang Nucleosynthesis (BBN) (see e.g. Tegmark & Zaldarriaga 2000; Jaffe et al. 2000). We note however that this discrepancy with the standard model is no more than 2σ when the most recent CMB anisotropy results are included Mason et al. 2001.

In recent months, several solutions have been advocated to reconcile BBN with the larger values of the baryon density suggested by the CMB data. In a recent letter by McGaugh (2000), the simple and controversial solution of considering only baryonic dark matter (BDM) plus a cosmological constant has been proposed. The lower CMB second peak can thereby be explained by the collisional damping expected in a BDM universe, while maintaining the baryon density at a level compatible with the BBN

constraint. As is well known, purely baryonic adiabatic models are in severe contradiction with the measured amplitude and shape of galaxy and cluster correlations. However, there remain serious issues about the possibly scale-dependent bias parameter. Hence McGaugh's result is interesting for various reasons: it resolves the disagreement with BBN by using a simpler model ($\Omega_{cdm} = \Omega_{hdm} = 0$), it demonstrates the model-dependence of theoretical fits to the CMB data by exploring a previously uncovered corner of parameter space and, finally, it questions our theoretical understanding of the structure formation process.

If it turns out that the CDM hypothesis does not work well at some cosmological epochs and scales, then even the exotic scenario of a purely baryonic model with modifications to standard gravity could become competitive, especially in view of the insistence of MOND proponents that this alternative theory of gravity merits serious examination (McGaugh & de Block 1998; Sanders 2000). Thus, it is timely to study how compatible the CMB data is with a BDM adiabatic universe, what consequences this assumption can have on the remaining cosmological parameters, and the need for CDM from independent observations of galaxy clustering.

In this *Letter*, we perform an analysis in cosmological parameter space. Using the recent CMB data *alone* we show that a purely baryonic adiabatic model appears unlikely if the universe is flat ($\Omega = 1$). By further combining these results with supernova type Ia (SNIa) data, we rule out baryonic adiabatic models with high confidence.

2. METHOD

The structure of the C_ℓ spectrum depends essentially on 3 cosmological parameters (see e.g. Hu, Sugiyama & Silk 1998; Efstathiou & Bond 1999; Melchiorri & Griffiths 2000 and references therein): the physical baryonic density $\omega_b = \Omega_b h^2$ and the overall matter density $\omega_m = \Omega_m h^2 = (\Omega_{cdm} + \Omega_b) h^2$ which define the size of the acoustic horizon at decoupling, and the 'shift' parameter R related to the geometry of the universe through $R = 2\sqrt{|\Omega_k|/\Omega_m}/\chi(y)$, with $y = \sqrt{|\Omega_k|} \int_0^{z_{dec}} [\Omega_m(1+z)^3 + \Omega_k(1+z)^2 + \Omega_\Lambda]^{-1/2} dz$, where the function $\chi(y)$ depends on the curvature of the universe and is y , $\sin(y)$ or $\sinh(y)$ for flat, closed and open models respectively.

A decrease in ω_{cdm} , with ω_b and R remaining constant, will reduce ω_m . This decreases the redshift of equality causing the peak in the spectrum to be shifted towards smaller angular scales. Constant ω_m can be maintained by increasing ω_b . However, an increase in ω_b will decrease the sound speed at decoupling $c_s \sim 1/\sqrt{3(1+3\omega_b/(4\omega_{rad}(1+z_{dec})))}$ again reducing the acoustic horizon size and shifting the peak to higher ℓ 's.

To keep the position of the first peak fixed while ω_b and ω_m are varied, R has to be appropriately tuned. This can be achieved by increasing Ω_Λ . The shift parameter R is sensitive to variations in Ω_Λ as $\Omega_m \rightarrow 0$, and for $\Omega_m = \Omega_b \sim 0.05$ a 10% increase in Ω_Λ can produce a 50% decrease in ℓ_{peak} . Thus, a purely baryonic model can produce the observed peak structure at $\ell \sim 200$ only if R is increased with respect to the corresponding flat CDM model. We can therefore anticipate that viable purely baryonic models will be Λ -dominated with closed geometries.

Motivated by these considerations, we compare recent CMB and SNIa observations with a set of models with parameters sampled as follows: $\Omega_m = \Omega_b = 0.015, \dots, 0.5$; $\Omega_\Lambda = 0.80, \dots, 1.04$ and $h = 0.40, \dots, 0.95$. We vary the spectral index of the primordial density perturbations within the range $n_s = 0.50, \dots, 1.50$ and we rescale the fluctuation amplitude by a pre-factor C_{10} , in units of C_{10}^{COBE} . The theoretical models are computed using the publicly available CMBFAST program (Seljak & Zaldarriaga 1996) and are compared with the recent BOOMERanG-98 and MAXIMA-1 results. The power spectra from these experiments were estimated in 12 and 10 bins respectively, spanning the range $25 \leq \ell \leq 785$. In each bin, the spectrum is assigned a flat shape, $\ell(\ell+1)C_\ell/2\pi = C_B$.

Following Bond, Jaffe & Knox (1999) we approximate the signal C_B inside the bin to be an offset lognormal distribution, such that the quantity $D_B = \ln(C_B + x_B)$ (where x_B is the offset correction) is a Gaussian variable. The likelihood for a given cosmological model is then defined by $-2\ln L = (D_B^{th} - D_B^{ex})M_{BB'}(D_B^{th} - D_B^{ex})$ where C_B^{th} (C_B^{ex}) is the theoretical (experimental) band power, x_B is the offset correction and $M_{BB'}$ is the Gaussian curvature of the likelihood matrix at the peak. We consider 10% and 4% Gaussian distributed calibration errors for the BOOMERanG-98 and MAXIMA-1 experiments respectively. We also include the COBE data using Lloyd Knox's RADPACK packages. Proceeding as in Dodelson & Knox (2000), we attribute a likelihood to a point in the $(\Omega_m = \Omega_b, \Omega_\Lambda)$ and (Ω_b, h) planes by finding the remaining 3 parameters that maximise it. We then define our 68%, 95% and 99% contours to be where the likelihood falls to 0.32, 0.05 and 0.01 of its peak value, as would be the case for a two dimensional multivariate Gaussian.

3. RESULTS

In the top panel of Figure 1, we plot likelihood contours in the $(\Omega_m = \Omega_b, \Omega_\Lambda)$ plane by applying the maximization/marginalization algorithm described above and using only the BOOMERanG-98 data and limiting our analysis to models with age $t_0 > 10\text{Gyr}$. As expected, the low sound speed at decoupling due to the high baryon content makes only closed models compatible with the observations. The deviation from flatness becomes less and less important as $\Omega_b \rightarrow 0$, as one would expect from the expression for R . However, the decrease in the redshift at equality causes positive curvature models to be preferred.

In the parameter range we are sampling, we find that flat models are excluded at $\sim 2\sigma$. In order to further test the discrepancy with a flat universe we also include CDM models in the analysis with $\Omega_{cdm} = 0.05, \dots, 0.45$. Comparing our models to the BOOMERanG-98 data alone and restricting the analysis to $\Omega = 1$, we find that $h^2\Omega_{cdm} > 0.04$ at the 95% C.L. Thus flat BDM models are excluded with significance.

For the BOOMERanG-98 analysis with $\Omega_{cdm} = 0$, the best-fitting model is a closed model with $\Omega_B = 0.135$, $\Omega_\Lambda = 1.0$, $n_s = 0.94$, $h = 0.45$, $C_{10} = 0.5$ and the BOOMERanG-98 calibration left untouched. This provides a good fit to the data (see Figure 2, top panel). Including the MAXIMA-1 data points produces a different best-fitting model: $\Omega_B = 0.035$, $\Omega_\Lambda = 1.00$, $n_s = 0.92$, $h = 0.85$, $C_{10} = 0.5$ with a $\sim 12\%$ upward calibration




FIG. 1.— The likelihood contours in the $(\Omega_b, \Omega_\Lambda)$ plane, with the remaining parameters taking their best-fitting values. Only the BOOMERanG-98 data is considered here. The contours correspond to 0.32, 0.05 and 0.01 of the peak value of the likelihood, which are the 68%, 95% and 99% confidence levels respectively. The dashed line going from left to right diagonally across the plot indicates flat models. The top panel is with a prior on the age of the universe $t > 10$ Gyr. The bottom panel includes the BBN prior $\Omega_b h^2 = 0.019 \pm 0.002$ and the SNIa data. The results of these two priors are incompatible at $\sim 3\sigma$.

for the BOOMERanG-98 points and a $\sim 5\%$ downward calibration for MAXIMA-1.

As the low value of C_{10} already suggests, these best-fitting models are in contradiction with the COBE data. Including COBE in the BOOMERanG-98 analysis does not significantly affect the best-fitting cosmological parameters which remain $\Omega_b = 0.135$, $\Omega_\Lambda = 1.0$, $n_s = 0.94$, $h = 0.45$, but the model now requires $C_{10} = 0.8$ and an upward shift in the calibration for BOOMERanG-98 of $\sim 30\%$. This model is ruled out by the COBE and BOOMERanG-98 data at the 95% C.L., although this discrepancy can be resolved by adding a gravitational wave (GW) contribution on large scales. The top panel of Figure 2 indicates that the inclusion of a GW component such that $C_{10}^T = C_{10}^S$ leaves the small scale behaviour practically unchanged but provides enough CMB power on large scales to match the COBE normalization.

A GW contribution as large as $C_2^T/C_2^S \sim 1$ is incompatible at the 95% C.L. with flat Λ CDM models when the BBN prior is assumed in CMB analyses (Kinney, Melchiorri & Riotto 2000). GWs leave a characteristic imprint on polarization power spectra, via fluctuations in the B magnetic-type-parities channel (Kamionkowski, Kosowsky & Stebbins 1996, Seljak & Zaldarriaga 1996) that vanish in the case of scalar fluctuations. However, the expected polarization amplitude for BDM models is very low, below $0.5\mu K$, and future experiments will not have the sensitivity to detect the B channel signal unless a substantial amount of reionization occurred in the past (Seljak & Zaldarriaga 1996).




FIG. 2.— In the top panel, the BOOMERanG-98 and the COBE data are plotted together with the best-fitting baryonic model to the BOOMERanG-98 data. In order to fit the COBE data, a tensor component must be added. For comparison, the Λ -CDM concordance model is also plotted. The middle panel shows the expected polarization signal of the best-fitting model. Including the tensor components produces B modes in the polarization spectra. In the bottom panel it can be seen that the best-fitting model to the BOOMERanG-98 data is in contradiction with the matter power spectrum measured by the PSCz survey.

One can further combine the CMB constraints with those obtained from the luminosity-distance measurements of high- z supernovae (Perlmutter et al. 1998, Schmidt et al. 1998) as we do in the bottom panel of Figure 1. The inclusion of the SNIa data does not greatly affect the constraints on Ω_Λ but, as expected, it rules out any models with low values of Ω_b . We find that $\Omega_b \geq 0.13$ at the 95% C.L. and an increasing inconsistency of BDM models with a flat universe.

Also plotted in the bottom panel of Figure 1 are the constraints obtained including a prior $\Omega_b h^2 = 0.019 \pm 0.002$ (Burles, Nollett & Turner 2000) inferred by measurements of primordial elements and assuming standard BBN. The results of the 2 priors are incompatible at more than 3σ .

It is interesting to further explore the compatibility of our results with independent observations by considering likelihood contours in the (Ω_b, h) plane. Figure 3 plots the results of our analysis with no priors, the results including the SNIa data and the results




FIG. 3.— The likelihood contours in the (Ω_b, h) , with the remaining parameters taking their best-fitting values to the BOOMERanG-98 data. The contours correspond to 0.32, 0.05 and 0.01 of the peak value of the likelihood, which are the 68%, 95% and 99% C.L. respectively. The filled contours correspond to an analysis with a $t > 10$ Gyr prior. The dashed, dotted and solid contours to the right of the plot are the result when the SNIa data is included. The bold black line to the left of the plot is the 95% C.L. region from BBN. with the BBN constraint.

We can see from this figure that purely baryonic models are in quite good agreement with BOOMERanG-98 and with the BBN constraints. However, when the SNIa data is included we are restricted to models with values of Ω_b that are too high and values of h that are too low to be consistent at $< 3\sigma$ with both the BBN constraint and the recent HST result $h = 0.72 \pm 0.08$ (Freedman et al. 2000).

Quite independently of the prior assumed, we find that the age of the universe in this scenario is constrained to be $t_0 = 24 \pm 2$ Gyr, nearly double that expected from standard CDM. The age of the universe must exceed the ages of the oldest globular clusters $t_{GC} = 14 \pm 2$ Gyr by an amount $\Delta t \sim 0.5 - 2$ Gyr, so the age of the universe required by these models seems to be too high unless globular cluster formation is delayed for a very long time.

Collisional damping erases fluctuations in the matter power spectrum on scales $< 100h^{-1}\text{Mpc}$. The predicted matter power spectrum from the best-fitting model (see Figure 2, bottom panel) is in contradiction with the decorrelated linear power spectrum extracted from the recent PSCz catalogue (Hamilton & Tegmark 2000). Allowing for a shift in the overall amplitude with a bias factor b where $P(k)_{PSCz} = b^2 P(k)$, we obtain a best-fitting bias $b = 4.5$ with a $\chi^2 = 155$ which is an extremely poor fit to 22 data points with one free parameter. The theoretical variance in baryonic matter fluctuations over a sphere of size $8h^{-1}\text{Mpc}$, for example, is $\sigma_8 \sim 10^{-2}$ to be compared with the observational value $\sigma_8 = 0.56\Omega_m^{-0.47} \sim 1.4$ (Viana & Liddle 1999). So, rescaling the galaxy data with a massive bias parameter may not be physically meaningful since no galaxies would have formed in order to allow there to be a predicted galaxy power spectrum.

Various phenomenological mechanisms can be proposed in an attempt to resolve the discrepancy with the matter power spectrum and render BDM models viable. One such possibility is to increase the spectral index of the primordial density fluctuations, n_s . However, the 'blue' models that are able to produce $\sigma_8 \sim 0.5$ require $n_s \geq 2$ which is outside the range $0.7 \lesssim n_s \lesssim 1.2$ allowed by infla-

tion (Steinhardt 1995). Also, such a high tilt will cause the CMB spectrum to become incompatible with observations.

Another option that could increase small scale density perturbations whilst leaving the CMB spectrum unchanged is to have a spectral index that varies with scale, $dn_s/d\ln k \neq 0$, as predicted by some inflationary models (Kosowsky & Turner 1995). Unfortunately, the variation needed, $n_s \gtrsim 1$, is much greater than that predicted by the most viable models.

More recently, there has been some investigation into features in the primordial power spectrum generated during inflation (Griffiths, Silk & Zaroubi 2000; ; Knebe, Islam & Silk 2001). It is possible that such features could increase the perturbations on small scales without affecting the CMB spectrum, although it is doubtful that a single feature would be able to resolve the observed discrepancy and such a mechanism is unlikely to be preferable to the assumption of CDM itself.

4. DISCUSSION

We have examined CMB anisotropies and large-scale structure observations in a purely baryonic dark matter universe. Our results suggest that a BDM adiabatic universe with a power-law primordial power spectrum can only reproduce the sub-degree CMB measurements if the universe is closed. Although the premise that $\Omega = 1$ is considered to be one of the basic predictions of the inflationary scenario, it is possible to construct inflationary scenarios that predict closed universes (see e.g. Linde 1995; Starobinsky 1996). It is interesting to note, however, that removing CDM immediately forces us to construct a more elaborate inflationary model, or, in other words, the most simple model needs non-baryonic matter to work.

When flatness was assumed, we found that $\omega_{cdm} > 0.04$ at 2σ with $L(\omega_{cdm} = 0)/L(\omega_{cdm} = 0.1) \sim 10^{-3}$. Therefore, the flatness constraint with BBN renders ΛBDM models less consistent with the CMB observations than the standard ΛCDM model. Also, ΛBDM models that fit the BBN constraints are degenerate with a non-physical region of the parameter space (with very high baryon content and small h). When any cosmological prior other than BBN is assumed in the analysis, the unphysical region turns out to be preferred. Including the SNIa constraints (or $\Omega_m = \Omega_b > 0.25$), for example, makes the parameter space incompatible at $> 95\%$ confidence with both the HST constraint $h = 0.72 \pm 0.08$ and with BBN.

As expected, all the BDM adiabatic models fail to reproduce both the CMB observations and the observed amount of galaxy clustering on $8h^{-1}\text{Mpc}$ scales. In order to solve all the discrepancies, that are not present in the standard ΛCDM scenario, we need to introduce *ad hoc* mechanisms which are unlikely to be preferable to the assumption of CDM itself. Nonetheless, BDM models leave a set of characteristic imprints such as large scale B -mode polarization and no third acoustic peak in the anisotropy spectrum that will allow future experiments to further scrutinize this hypothesis.

We thank Pedro Ferreira, Arthur Kosowsky, Andrew Liddle, Francesco Melchiorri, Gary Steigman for useful conversations and the High-Z Supernova Search Team for providing the SNIa likelihoods.

REFERENCES

- Bahcall N. et al. 1999, *Science*, 284, 1481
 Barriga J., Gaztanaga E., Santos M., Sarkar S., 2000, *astro-ph/0011398*
 Bergstrom L., 2000, *Rep. Prog. Phys.*, 63, 793
 de Bernardis P. et al., 2000, *Nature*, 404, 955
 Binney, J. Gerhard, O. and Silk, J. 2001, *MNRAS*, 321, 471.
 Bond J. R., Jaffe A. H., Knox L. 1998, *Phys Rev D*, 57, 2117
 Burkert A., 2000, *astro-ph/0002409*
 Burles S., Nollett K. M., Turner M. S., 2000, *astro-ph/0010171*
 Dodelson, S., Knox, L., *astro-ph/9909454*
 Doroshkevich A. G., Zel'dovich Y. B., Sunyaev R. A., 1978, *Sov. Astronom.*, 21, 523
 Efstathiou G., Bond J. R., 1999, *MNRAS*, 304, 75
 Freedman W. L. et al., 2000, *astro-ph/0012376*
 Griffiths L. M., Silk J., Zaroubi S., 2000, *MNRAS accepted*, *astro-ph/0010571*
 Hamilton A. J. S., Tegmark M., 2000, *astro-ph/0008392*
 Hanany S. et al., 2000, *astro-ph/0005123*
 Hu W., Sugiyama N. & Silk J., 1998, *Nature*
 Jaffe A. H. et al., 2000, *astro-ph/0007333*
 Kaiser N., 1983, *MNRAS*, 202, 1169
 Kamionkowski M., Kosowsky A., Stebbins A., 1996, *Phys. Rev. Lett.*, 78, 2058
 Kinney W., Melchiorri A., Riotto A., 2000, *astro-ph/0007345*
 Knebe A., Islam R. R., Silk J., *astro-ph/0103082*
 Kosowsky A., Turner M. S., *Phys. Rev. D*, 1995, 52, 1739
 Linde A., 1995, *Phys. Lett.*, B, 351, 99
 Madsen J., 2000, *astro-ph/0006074*
 McGaugh S. S., de Block W. J. G., 1998, *ApJ*, 499, 66
 McGaugh S. S., 2000, *ApJ*, 541, L33
 Mason, B., Cartwright, J., Padin, S., Pearson, T., Readhead, A., Shepherd, M., Sievers, J. and Udomprasert, P. 2001, *astro-ph/0101171*.
 Melchiorri A., Griffiths L. M., 2000, *astro-ph/0011147*
 Moore B. et al., 2000, *ApJ*, 535, L21
 Perlmuter S. et al., *astro-ph/9812133*
 Sanders R. H., 2000, *astro-ph/0011443*
 Schmidt B. P. et al. 1998 *ApJ*, 507, 46
 Seljak U., Zaldarriaga M. 1996, *ApJ*, 469, 437
 Seljak U., Zaldarriaga M., 1996, *Phys. Rev. Lett.*, 78, 2054
 Sellwood J. A., Kosowsky A., 2000, *astro-ph/0009074*
 Spergel D. N., Steinhardt P. J., 1999, *astro-ph/9909386*
 Starobinsky A., 1996, *astro-ph/9603075*
 Steinhardt P., 1995, *astro-ph/9502024*
 Steinhardt P. J., Wang L., Zlatev I., 1999, *Phys. Rev. D* 59, 123504
 Tegmark M., Zaldarriaga M., 2000, *Phys. Rev. Lett.*, 85, 2240
 Viana P., Liddle A. R., 1999, *MNRAS*, 303, 535
 White S. D. M., Frenk C., Davis M., 1983, *ApJ*, 274, L1
 Wilson M. L., Silk J., 1981, *ApJ*, 243, 14
 Zwicky F., 1933, *Helv. Phys. Acta*, 6, 110